


OBSERVATION OF COSMIC RAYS ANISOTROPY ABOVE TEV ENERGIES IN ICECUBE

Simona Toscano *on behalf of the* **IceCube** collaboration



**3rd Roma International Conference
on Astro-particle Physics**

**25-27 May 2011
Roma Italy**



Outline

- * The **IceCube** detector
- * Energy dependence of the **large scale anisotropy** (*paper in preparation*):
 - ▶ preliminary results at 20 and 400 TeV.
 - ▶ solar dipole
- * **Medium and small scale** structures (submitted to ApJ, **arXiv:1105.2326**):
 - ▶ analysis
 - ▶ systematics
- * Conclusions



Bartol Inst, Univ of Delaware
Penn State
UW-Madison
UW-River Falls
LBNL, Berkeley
UC Berkeley
UC Irvine
Univ. of Alabama
Clark-Atlanta University
Univ. of Maryland
University of Kansas
Southern Univ. and A&M College
University of Alaska, Anchorage
Georgia Tech
Ohio State



Imperial College, London, UK
University of Oxford, UK



Université Libre de Bruxelles, Belgium
Vrije Universiteit Brussel, Belgium
Université de Mons-Hainaut, Belgium
Universiteit Gent, Belgium



Utrecht University, Netherlands



EPFL, Lausanne, Switzerland



Universität Mainz, Germany
DESY Zeuthen, Germany
Universität Wuppertal, Germany
Universität Dortmund, Germany
Humboldt Universität, Germany
MPI, Heidelberg
Ruhr-Universität, Bochum



Chiba University, Japan



University of West Indies
Barbados



Uppsala Universitet, Sweden
Stockholm Universitet, Sweden
Kalmar Universitet, Sweden



University of Canterbury,
Christchurch, New Zealand



IceCube Collaboration



10 countries
36 institutions
~260 collaborators

The IceCube detector


- ➔ **G. Sullivan** - *Status and Recent Results from the IceCube km³ Neutrino Detector* (tomorrow Plenary)
- ➔ **T. DeYoung** - *Particle physics in ice with IceCube DeepCore* (today next Parallel session)

- **86 strings**
- **5160 DOMs**
- **17 m vertical spacing**
- **125 m between strings**

Digital Optical Module




IceCube configurations



Construction finished on December 2010


IceCube configurations

IC-1
04-05 Season




Construction finished on December 2010

IceCube configurations




Construction finished on December 2010

IceCube configurations



Construction finished on December 2010

IceCube configurations




IC-40
07-08 Season

Construction finished on December 2010


IceCube configurations

IC-59
08-09 Season




Construction finished on December 2010

IceCube configurations



Construction finished on December 2010

IceCube configurations




Construction finished on December 2010

Data set

*IceCube collects billions of down-going muons generated in CR air showers





***32 billion** of events collected between May 2008/May 2009 with **20 TeV** median energy




IceCube-59	Level 1 "Muon Filter"	DST
Live Time	96%	96%
Trigger rate	22 Hz	1.4 kHz
N_{events}	8.0×10^8	3.2×10^{10}
Angular resolution	$< 1^\circ$	3°
Energy resolution	$\Delta \log_{10}(E) = 0.3$	$\Delta \log_{10}(E) = 0.5$

Observation of the CRs large scale anisotropy

There have been several observations of *large-scale, part-per-mille anisotropy* in cosmic ray arrival directions between 0.1 and 100 TeV.



Northern Sky




Southern Sky

Tibet ASy	M. Amenomori et al., Astrophys. J. 626 (2005) L29
SuperK	G. Guillian et al., Phys. Rev. D 75 (2007) 062003
Milagro	A. Abdo et al., Astrophys. J. 698 (2009) 2121
ARGO-YBJ	S. Vernetto, Proc. 31st ICRC, 2009
EAS-Top	M. Aglietta, Astrophys. J. 692 (2009) L130
IceCube	R. Abbasi et al., Astrophys. J. 718 (2010) L194

Large scale anisotropy

- * IceCube observed a large scale anisotropy at 10^{-3} level for the first time in the Southern Sky.
- * The anisotropy appears to be a continuation of large scale structures observed in the Northern Hemisphere.



Relative intensity of the cosmic ray event rate in equatorial coordinates: for each declination belt of width 3° , the plot shows the number of events relative to the average number of events in the belt.

Energy dependence of the anisotropy



300 TeV

Tibet - III

Amenomori et al., Science Vol. 314, pp. 439, 2006

Amplitude: $(3.2 \pm 2.6) \times 10^{-4}$

consistent with **no anisotropy**



110 TeV

EAS-Top

Aglietta et al., ApJ 692, L130, 2009



370 TeV

Amplitude (370 TeV): $(6.4 \pm 2.5) \times 10^{-4}$





consistent with **no anisotropy** if systematics were taken into account properly

Relative Intensity

Equatorial sky maps in HEALPix with $N_{\text{Side}} = 16$, $\text{pix resol} \sim 3^\circ$




Preliminary






IO

Earth's motion around the Sun: the Solar dipole



$$\frac{\Delta I}{I} = (\gamma + 2) \frac{v}{c} \cos \theta$$





$v = 29.8 \pm 0.5 \text{ km/s}$



Energy dependence of the Solar dipole

- * IceCube observes the Solar dipole in both energy bins. The observed amplitude is compatible with the expectations within the stat. and sys. uncertainties.
- * The observation of the solar dipole supports the observation of the sidereal anisotropy in cosmic ray arrival direction.

relative intensity Vs. $(\alpha[^\circ] - \alpha_{\text{SUN}}[^\circ])$




Preliminary

Small scale anisotropy

Several experiments have discovered anisotropies on scales of about 10°


- * Milagro observes two localized regions with **significance $> 10\sigma$** in the total data set of $2.2 \cdot 10^{11}$ events recorded over 7 years. The “hot” regions have fractional excesses of order several times **10^{-4}** relative to the background.
- * Same structures observed by ARGO-YBJ.



A. Abdo et al., PRL 101 (2008) 221101

Milagro

Median Energy: 1 TeV




S. Vernetto, Proc. 31st ICRC, 2009

ARGO-YBJ

Median Energy: 2 TeV

Relative Intensity map

Equatorial sky maps in HEALPix: equal area pixel (size ~ 0.9°)



Sky map created using the background estimation technique from real data:

- N_i : number of data events in the i^{th} pixel.
- $\langle N_i \rangle$: expected number of events in an isotropic sky (time scrambling in 24 hr) in the i^{th} pixel.
- Relative Intensity:

$$\frac{\Delta N_i}{\langle N \rangle_i} = \frac{N_i(\alpha, \delta) - \langle N_i(\alpha, \delta) \rangle}{\langle N_i(\alpha, \delta) \rangle}.$$



Relative intensity map is *not isotropic*. In IceCube-59, the *strong large scale structure* already observed in IceCube-22 data is visible in the “raw” data.

Significance map

Significance calculation:


$$s = \sqrt{2} \left\{ N_{\text{on}} \ln \left[\frac{1 + \alpha}{\alpha} \left(\frac{N_{\text{on}}}{N_{\text{on}} + N_{\text{off}}} \right) \right] + N_{\text{off}} \ln \left[(1 + \alpha) \left(\frac{N_{\text{off}}}{N_{\text{on}} + N_{\text{off}}} \right) \right] \right\}^{1/2} \quad \alpha = 1/20$$

Li, T., & Ma, Y. 1983, ApJ, 272, 317



Power spectrum


Angular size $\theta \sim \frac{180^\circ}{\ell}$



Multipole expansion: $\delta I(\mathbf{u}_i) = \sum_{\ell=1}^{\infty} \sum_{m=-\ell}^{\ell} a_{\ell m} Y_{\ell m}(\mathbf{u}_i) \quad \mathcal{C}_{\ell} = \frac{1}{2\ell + 1} \sum_m |a_{\ell m}|^2$

Power spectrum

Angular size $\theta \sim \frac{180^\circ}{\ell}$



Multipole expansion: $\delta I(\mathbf{u}_i) = \sum_{\ell=1}^{\infty} \sum_{m=-\ell}^{\ell} a_{\ell m} Y_{\ell m}(\mathbf{u}_i)$ $\mathcal{C}_{\ell} = \frac{1}{2\ell + 1} \sum_m |a_{\ell m}|^2$

Dipole and quadrupole fit

$$\delta I(\alpha, \delta) = m_0$$

$$+ p_x \cos \delta \cos \alpha + p_y \cos \delta \sin \alpha + p_z \sin \delta$$

$$+ \frac{1}{2} Q_1 (3 \cos^2 \delta - 1) + Q_2 \sin 2\delta \cos \alpha + Q_3 \sin 2\delta \sin \alpha + Q_4 \cos^2 \delta \cos 2\alpha + Q_5 \cos^2 \delta \sin 2\alpha$$

monopole


dipole

quadrupole


Coefficient	Fit Value
m_0	0.320 ± 2.264
p_x	2.435 ± 0.707
p_y	-3.856 ± 0.707
p_z	0.548 ± 3.872
Q_1	0.233 ± 1.702
Q_2	-2.949 ± 0.494
Q_3	-8.797 ± 0.494
Q_4	-2.148 ± 0.200
Q_5	-5.268 ± 0.200

$$\chi^2/\text{ndf} = 14743.4/14187$$

$$\text{Pr}(\chi^2|\text{ndf}) = 5.5 \times 10^{-4}$$



Residual map




No structures seem to be present: we need to smooth the map.

MAP SMOOTHING SCAN

Scan from 1 - 30° in smoothing
Different regions have different optimal angular smoothing
Significances are pre-trial



MAP SMOOTHING SCAN

Scan from 1 - 30° in smoothing
Different regions have different optimal angular smoothing
Significances are pre-trial





Identification of significant structures

region	right ascension	declination	optimal scale	peak significance	post-trials
1	$(122.4^{+4.1}_{-4.7})^\circ$	$(-47.4^{+7.5}_{-3.2})^\circ$	22°	7.0σ	5.3σ
2	$(263.0^{+3.7}_{-3.8})^\circ$	$(-44.1^{+5.3}_{-5.1})^\circ$	13°	6.7σ	4.9σ
3	$(201.6^{+6.0}_{-1.1})^\circ$	$(-37.0^{+2.2}_{-1.9})^\circ$	11°	6.3σ	4.4σ
4	$(332.4^{+9.5}_{-7.1})^\circ$	$(-70.0^{+4.2}_{-7.6})^\circ$	12°	6.2σ	4.2σ
5	$(217.7^{+10.2}_{-7.8})^\circ$	$(-70.0^{+3.6}_{-2.3})^\circ$	12°	-6.4σ	-4.5σ
6	$(77.6^{+3.9}_{-8.4})^\circ$	$(-31.9^{+3.2}_{-8.6})^\circ$	13°	-6.1σ	-4.1σ
7	$(308.2^{+4.8}_{-7.7})^\circ$	$(-34.5^{+9.6}_{-6.9})^\circ$	20°	-6.1σ	-4.1σ
8	$(166.5^{+4.5}_{-5.7})^\circ$	$(-37.2^{+5.0}_{-5.7})^\circ$	12°	-6.0σ	-4.0σ



Systematics: Solar Dipole

We are sensitive to the motion of the Earth around the Sun (10^{-4} effect is expected): visible when UT is used in local-celestial coord. transformation.




Coefficient	Fit Value ($\times 10^{-4}$)
m_0	-0.029 ± 0.058
p_x	0.017 ± 0.142
p_y	-3.661 ± 0.142
p_z	-0.027 ± 0.072

$$\chi^2/\text{ndf} = 14206.8/14192$$




$$\text{Pr}(\chi^2|\text{ndf}) = 0.416$$

Systematics: previous data sets







Different geometries, same structure
Signal grows with statistics

Systematics: previous data sets








Different geometries, same structure
Signal grows with statistics

Systematics: previous data sets



Different geometries, same structure
Signal grows with statistics


Systematics: previous data sets



Different geometries, same structure
Signal grows with statistics

Milagro + IceCube combined map

IceCube map contains all data from IC22, IC40 and IC59 data sets



Milagro map:

[Abdo, A. A., et al. 2008, Phys. Rev. Lett., 101, 221101]

- 2.2×10^{11} events
- time scrambling (2 hr)
- 10° smoothing
- median energy **1 TeV**

IceCube map:

- 5.6×10^{10} events
- time scrambling (4 hr)
- 10° smoothing
- median energy **20 TeV**



Conclusions

- * **IceCube** detector **completed in December 2010** and now taking data in its final configuration (86 strings).
- * **Large scale anisotropy:**
 - ▶ First observation of sidereal anisotropy @ 400 TeV in southern hemisphere.
 - ▶ Sidereal anisotropy at 20 TeV confirms previous observation.
 - ▶ Indication of a persistence of anisotropy @ 400 TeV: evidence of a “dip”.
- * **Small and medium scale** structures:
 - ▶ Southern sky in TeV cosmic rays shows significant anisotropy across a wide range of angular scales (10-180 degrees).
 - ▶ Features similar to what observed in the Northern Sky

BACKUP SLIDES

Energy estimation

