

# Technical progress

- detector performance, calibration, R&D efforts

Albrecht Karle

March 2019

# The IceCube Neutrino Observatory

**IceTop** (surface array): 81 stations

**IceCube**: 86 strings


5160 optical sensors over 1 km<sup>3</sup> volume

17 m vertical spacing


125 m horizontal spacing

Highly stable operation.

Since 2016: **livedtime** > 99.5%



**DeepCore** (low energy threshold)





**South Pole 10m Telescope**



**MAPO**

**TOS - Drilling site (79 & 80 in 10/11)**





**IceCube Laboratory (ICL)**

**IceCube Enhanced Hot Water Drill (EHWD)**




3100 sensors are deployed  
to a depth between 1500  
and 2500m.






# Detector Uptime



# PMT gain stability 2011 - 2016

No indication for any changes since 2016.




DOM gain appears stable!

(PMT gain of  $1E7$  is small.  
Noise rates are small.  
→ Very small integrated current on anode.  
→ No aging from that.


# Time resolution: ~1ns for bright pulses

- Time difference between neighboring DOMs fired with (bright) flasher pulses: ~1 ns.  
(this includes clock timing)




Single photoelectron pulse resolution limited by PMT.  
RMS in the peak: ~2ns

Lab measurement with laser.



# Understanding the ice

## 1. Vertical structure of ice parameters




Scattering (eff.): 20 – 50 m  
Absorption: 100 – 200 m

Measurement of South Pole ice transparency with the IceCube LED calibration system,


Aartsen et al., (IceCube Coll.), NIMA55353  
<http://arxiv.org/abs/1301.5361>

## 2. Azimuthal variation in of scattering


Less scattering in direction of ice flow:  
→ up to ~10% /100m variation in amplitude



## 3. Ice layers are tilted – not planar



## 3. Scattering function







# Systematic uncertainties: DOM and local ice

We plan to map the full surface sensitivity of every DOM precisely cable position to  $<3^\circ$  (can be determined with local LEDs), then fit effect of hole ice.

Current picture of hole ice



# DOM and local ice


Images taken with camera ("Swedish Camera") during refreeze process:



Hole ice visible on the right.  
Need to determine the effect  
for every single DOM.


# Cable shadow

Cable diameter: 4.5cm




DOM sphere: 32.5

PMT cathode diameter: 22 cm





Azimuthal DOM response:  
Simulated effect on receiving  
DOM from flashers at close  
distance.




# Built-in inclinometers vs DOM tilt fit

## Indication of real tilt for 2 DOMs (out of 48)!

4 dozen DOMs have a built-in inclinometer, mounted on the mainboard, most of them have measured very small tilts, while 2 have tilts in excess of 20 degrees.




Example of DOM level calibration work:  
determined position of individual cables near DOM to few degree  
precision



# Types of events and interactions

**Charged-current  $\nu_\mu$**




Up-going (throughgoing) track

Factor of  $\sim 2$  energy resolution  
 $\sim 0.5^\circ$  angular resolution

**$0.3^\circ$  above 100 TeV**

**Neutral-current /  $\nu_e$**




Isolated energy deposition  
 (cascade) with no track

15% deposited energy resolution  
 10-15° angular resolution (above 100 TeV)  
 Working on improving that.



**Charged-current  $\nu_\tau$**

(simulation)



“Double-bang”

(none observed yet:  $\tau$   
 decay length is 50 m/  
 PeV)


ID: above  $\sim 100$  TeV  
 (two methods)

# Bright DOMs


---

DOMs with  $Q_{\text{bright}} > 10 * Q_{\text{avg}}$  are classified as "Bright"


PMT not necessarily saturated, but excluded because unmodeled systematic uncertainties start to dominate at high photon statistics




# Observation of a 6 PeV neutrino



Resonance:  $E_\nu = 6.3$  PeV  
Typical visible energy is 93%



**Work in progress**



Event identified in a partially-contained PeV search (PEPE)

Deposited energy:  $5.9 \pm 0.18$  PeV (stat only)

ICRC 2017 [arXiv:1710.01191](https://arxiv.org/abs/1710.01191)




# A neutrino event near Glashow resonance

Interesting event found in expanded search.

Charge: 200,000 photoelectrons

**Energy:  $\sim 6$  PeV**



Ref: ICRC 2017, L. Lu (IceCube C.)

# Energy resolution is critical






Figure from Tianlyu that shows difference of resolution with and without systematic errors.




# Local effects: DOM orientation and cable position

Without local effects




With local effects




# Higher level performance parameters

## Angular resolution for muon neutrinos

2019



2011



# Moon shadow


Angular resolution for muon neutrinos

# Simulated cascade

Bright DOMs:  $Q > 10 * Q_{avg}$

DirectFit (Dima) reconstructing data events with direct photon simulation with ppc.

PeV-cascade: (Sim: 3.2, Reco: Mie)




Angular resolution improves with increased values of total charge used and maximum charge per DOM




-> see Tianlu Yuan talk

Starting muon  
"Dr. Strangepork"  
contained vertex,  
Deposited energy: 7






# Milestone 2013: Discovery of diffuse cosmic neutrino flux






Initially with 2 - 3 years of data



Contained vertex events  
7 years of data


Through and upgoing muons  
8 years of data



# R&D

- R&D related to M&O and continued optimization of IceCube proper
  - Surface instrumentation
  - SpiceCore
- R&D geared towards the future: Upgrade and Gen2
  - Detector R&D, new optical modules

# Science case



## **Snow attenuation mitigation:**



- Measure the effect of snow on IceTop tank sensitivity, binned by energy, zenith, and radial distance from shower core.
- Recover the sensitivity to low-energy showers that are currently not detected by tanks buried under several feet of snow.


## **Veto efficiency improvement:**

- By adding scintillators with a similar coverage as IceTop, the energy threshold at which the veto becomes efficient at a  $10^4$  to  $10^5$  rejection factor is estimated to be lower by a factor of two.

## **R&D for future detector upgrades:**



- A new, scalable precision timing and high-speed communications scheme for IceCube M&O and possible future projects.
- Efficient trenching procedures for instrumentation installation.
- Mechanical solutions to raise scintillator panels above the snow during the period of array deployment.






→ Position Y (m)

→ Position X (m)




# Exploring Southern sky with using veto techniques

Event near Galactic Center



Expect 20 to 100 events in southern sky in 10 years depending on spectrum.




Highest Energy Event

Self veto optimized for starting muon tracks.  
High purity astrophysical  $\nu_{\mu}$  events at  $\sim 10$  TeV!

# Multimessenger astronomy in real time - flares

## Implementation of efficient realtime system online

Technical progress:  
TXS alert published 43  
seconds after interaction.




# IceCube Upgrade (a step towards Gen2)

Funded.

7 strings in center of IceCube,  
densely instrumented

Science goals:

- $\nu_\mu$  disappearance
- $\nu_\tau$  appearance
- Precise calibration of IceCube optical properties and DOM response



A big step towards IceCube-Gen2



# IceCube-Gen2

The next Generation IceCube: from discovery to astronomy

## Multi-component observatory:


- IceCube-Gen2 High-Energy Array
- Surface air shower detector
- Sub-surface radio detector

Surface Area:  $\sim 6.5 \text{ km}^2$  (0.9)

Instrumented depth: 1.26 km (1.0)


Instrumented Volume:  $8 \text{ km}^3$

Order of magnitude increase  
of contained event rate at high  
energies.





Artist conception  
Here: 120 strings at 300 m spacing

# IceCube Gen2 schedule




*Funded*



*pending*





Thank you!






New event selections “below”  
HESE and throughgoing muons...

# From High to Medium energy: Part 1 - MESE

Follow-up analysis to [arxiv.org/1410.1749](https://arxiv.org/abs/1410.1749)

- 2 years  $\rightarrow$  7 years
- and optimized



# From High to Medium energy:

Low-threshold

7-yr unfolding

- Unfolding to  $\nu_e$ 
  - assume isotropic
  - compatible with tl

